Alpha (α) thalassaemia

What is alpha (α) thalassaemia?

Thalassaemia is a group of blood disorders affecting haemoglobin production. Haemoglobin is a protein in the blood that carries oxygen around our bodies.

Thalassaemia is passed from parent to child in genes. Genes carry information about human characteristics such as eye colour, hair colour and haemoglobin.

Thalassaemia is inherited.

Thalassaemia is <u>not</u> contagious.

Thalassaemia is <u>not</u> transmitted by germs.

Sometimes changes occur to genes, resulting in medical conditions. Such changes occur to alpha globin genes in alpha (α) thalassaemia:

- A person normally inherits four α globin genes for the production of the alpha globin protein in haemoglobin.
- A person may have two or three of the normal four alpha globin genes for haemoglobin production. This person is called a carrier of a thalassaemia and is <u>healthy</u>.
- Carriers may be at risk of having a child affected with Haemoglobin H disease or Bart's hydrops fetalis if their partner is also a carrier of certain types of α thalassaemia.
- When a person has only one alpha globin gene, they have Haemoglobin H disease and require regular medical care. Individuals with Haemoglobin H disease may experience lifelong anaemia of mild to moderate degree. Occasionally it may be severe.
- When a person has no alpha globin genes, they have a severe condition called **Bart's hydrops** fetalis. Bart's hydrops fetalis affects a foetus long before birth, resulting in death during pregnancy or shortly after birth. This is a fatal condition which is dangerous for both the mother and baby during pregnancy.

Treatment

Those with Haemoglobin H disease may require blood transfusions to correct anaemia. There is no treatment or cure for Bart's hydrops fetalis.

The health of carriers of $\boldsymbol{\alpha}$ thalassaemia

A carrier can expect to be <u>healthy</u>. It is important that their doctor knows they are a carrier of a thalassaemia.

Alpha thalassaemia and family planning

The genes for α thalassaemia are common in people of Asian origin, as well as those of African, Middle Eastern and Mediterranean origin.

Couples planning a family, or early in pregnancy, should have a blood test to determine whether or not they are carriers, if the origin of either of their families is one of the areas listed above; or if they have a family history of any blood disorder or anaemia. This test is needed to determine if there is any risk of having a child affected by a genetic blood disorder.

Those at risk of having an affected child have options. These conditions can be diagnosed as early as the 12th week of pregnancy. Termination of pregnancy can then be considered, if appropriate. People can adopt or can consider assisted reproductive techniques (such as preimplantation genetic diagnosis, the use of donor eggs or donor sperm). Others may choose to take the chance of having an affected child. All of these options can be discussed with a Genetic Counsellor.

Testing can be arranged by your local doctor or by contacting the hospitals listed at the end of this pamphlet.

Important information for your family

If you are a carrier of alpha thalassaemia, other members of your family may also be carriers and at risk of having children with a severe form of alpha thalassaemia. It is recommended that other family members **and** their partners be tested for their carrier status **prior** to having children of their own. DNA testing is utilized to detect carriers of alpha thalassaemia.

Figure 1:

Carrier parents have 2 alpha (α) globin genes



With each pregnancy, this couple has a:

- 1 in 4 chance of having a child who is not a carrier of α thalassaemia.
- 2 in 4 chance of having a healthy carrier.
- 1 in 4 chance of Bart's hydrops fetalis.

Figure 2:

Carrier parents have 2 and 3 α globin genes



With each pregnancy, this couple has a:

- 1 in 4 chance of having a child who is not a carrier of α thalassaemia.
- 2 in 4 chance of having a healthy carrier.
- 1 in 4 chance of having a child with Haemoglobin H disease.

Figure 3:

Carrier parents have 2 α globin genes



With **<u>each</u>** pregnancy, this couple has a:

- 2 in 4 chance of having a healthy carrier.
- 2 in 4 chance of having a child with Haemoglobin H disease.

Useful contacts

Monash Medical Centre - Medical Infusion Unit (State Thalassaemia Service) 246 Clayton Road, Clayton VIC 3168 Phone: +61 3 9594 2756 or +61 3 9594 3154

Royal Melbourne Hospital (5 West Day Medical Centre) 300 Grattan Street, Parkville VIC 3052 Phone: +61 3 9342 7092 or +61 3 9342 8835



Thalassaemia and Sickle Cell Australia Moorleigh Community Village 92-94 Bignell Road Bentleigh East VIC, 3165, Australia

Phone : (03) 7015 5637 Email: info@tasca.org.au Mail to: P.O. Box 3076 Moorabbin East VIC, 3189, Australia

This fact sheet has been translated into a number of different languages and they are available to download at *www.tasca.org.au* © Thalassaemia and Sickle Cell Australia 2020